




JEB™

p-ISSN: 0254-8704
e-ISSN: 2394-0379
CODEN: JEBIDP

Effect of nitrogen and phosphorus on growth and microcystin production in three *Microcystis* species



Authors Info

K.L. Lee¹, K.Y. Jung², J.H. Kim⁴,
Y.S. Kim³ and H.S. Kim^{3*}¹Watershed Ecology Research Team, National Institute of Environmental Research, Incheon, 22689, South Korea²Nakdong River Environment Research Center, National Institute of Environmental Research, Gyeongsangbuk-do, 40103, South Korea³Department of Biology, Kyungpook National University, Daegu, 41566, South Korea⁴Microorganism Resources Division, National Institute of Biological Resources, Incheon, 22689, South Korea*Corresponding Author Email : kimhsu@knu.ac.kr

Key words

Cyanobacterial blooms
Freshwater ecosystem
Microcystin
Microcystis species
Nitrogen & Phosphorus

Publication Info

Paper received : 08.04.2017
Revised received : 13.07.2017
Re-revised received : 06.09.2017
Accepted : 29.09.2017


Abstract

Aim : The effects of nitrogen and phosphorus concentration on growth and microcystin production were investigated in three species of bloom-forming *Microcystis* isolated from two South Korean freshwater systems.

Methodology : Three species of cyanobacteria were collected from Yeongchun Dam and Ankei Dam in Kyungpook Province, South Korea. Culture experiments were conducted at $25 \pm 1^\circ\text{C}$ under cool white fluorescent light (ca. $50 \mu\text{mol m}^{-2} \text{s}^{-1}$) in media with different concentrations of nitrogen (0 to 20 mg l^{-1}) and phosphorus (0 to 5 mg l^{-1}). Cell numbers were determined in a hemocytometer for calculation of growth rate. Microcystin was analysed using high pressure liquid chromatography.

Results : The highest growth rate (μ_{max}) and maximal microcystin production occurred at nitrogen concentrations of 10 and 20 mg l^{-1} in all three species. The response to phosphorus concentration was more complex. The highest growth rate (μ_{max}) of *M. aeruginosa*, *M. ichthyoblabe* and *M. viridis* occurred at phosphorus concentrations of 0.5 mg l^{-1} , 0.1 mg l^{-1} , and 3 mg l^{-1} , respectively. *M. aeruginosa* also had maximal microcystin production at 0.5 mg l^{-1} P. In contrast, *M. ichthyoblabe* and *M. viridis* had high microcystin production at 0 mg l^{-1} and at 5 mg l^{-1} P (the highest tested concentration), and low microcystin production at 0.1 mg l^{-1} P. Thus at 0.1 mg l^{-1} , *M. ichthyoblabe* had the highest growth rate but produced least amount of microcystin. The types of microcystins produced varied according to species and nutrient conditions.

Interpretation : Microcystin production and growth in *Microcystis* species isolated from South Korea varied according to species and nutrient conditions. These species responded similarly to different nitrogen concentrations, but differently to different phosphorus concentrations.



Introduction

Cyanobacterial blooms are common in water bodies all over the world, and cyanobacterial toxins are a serious threat to the health of aquatic animals and humans. Microcystins are hepatotoxins produced by certain freshwater cyanobacteria and have well-documented harmful effects in humans (Kuiper-Goodman et al., 1999; Falconer, 2001). *Microcystis*, *Anabaena*, *Oscillatoria*, *Nostoc* and *Anabaenopsis* can produce microcystins (Kaebernick and Neilan, 2001) and species of the genus *Microcystis* are the best-known microcystin-producing cyanobacteria (Park et al. 1998; Kurmayer et al., 2002; Via-Ordorika et al., 2004; Ozawa et al., 2005; Znachor et al., 2006). The genus *Microcystis* consists of 15 species, including *M. aeruginosa*, *M. ichthyoblabe*, *M. flos-aquae*, *M. novacekii*, *M. viridis* and *M. wesenbergii*, all of which have characteristic cell and colony morphologies (Komárek and Anagnostidis, 1999).

Microcystin production is species- and strain-specific (Oh et al., 2000; Rohlack et al., 2001; Kurmayer et al., 2002; Via-Ordorika et al., 2004; Yéprémian et al., 2007). Thus, different species and strains produce different type and amount of microcystins, depending on environmental conditions. Studies carried of microcystin production, have mainly focused on *Microcystis aeruginosa* (Codd and Poon, 1998; Watanabe, 1996; Oh et al., 2000). However, *M. ichthyoblabe* and *M. viridis* are also responsible for *Microcystis* blooms (Park et al., 1998; Sabour et al., 2002).

It is necessary to understand the physiological and ecological characteristics of microcystin production by different species of *Microcystis* to better ensure a safe supply of drinking water. Nevertheless, only few studies have compared microcystin production using cultures of different *Microcystis* species and there are no data on the growth of Korean strains of *Microcystis* in culture. In this study, the authors investigated the effects of nitrogen and phosphorus on the growth and microcystin production of three species of bloom-forming *Microcystis* (*M. aeruginosa*, *M. ichthyoblabe* and *M. viridis*) that were isolated from Korean freshwater systems and grown in culture.

Materials and Methods

Strain isolation : Three species of *Microcystis* (*M. aeruginosa* YC, *M. ichthyoblabe* AK and *M. viridis* AK) were isolated from Youngcheon dam and Ankei dam during cyanobacterial blooms. These dams are adjacent and linked by a number of waterways. The species composition of the cyanobacterial communities in two dams were similar at the time of blooms.

Culture conditions : All culture experiments were conducted using CB medium (Watanabe, 1996). Unialgal stock cultures were established and maintained in CB medium (buffered to pH 9.0 with carbon free NaOH) at $25 \pm 1^\circ\text{C}$ and a light intensity of approximately $50 \mu\text{mol m}^{-2} \text{s}^{-1}$ under continuous cool white fluorescent light. For nutrient experiments, clones selected from

the stock cultures at the exponential growth phase were first adapted to nutrient-depleted medium (no N or P) for one week, and were then inoculated (initial cell density: ca 5000 cells) into medium with different levels of nitrogen (0, 1, 3, 5, 10, 20 mg l⁻¹) and phosphorus (0, 0.1, 0.5, 1, 3, 5 mg l⁻¹). NO₃⁻ and PO₄³⁻ were the sources of nitrogen and phosphorus. These cultures were grown for two weeks as described above. Experiments were conducted in triplicate and cell numbers were determined with a hemocytometer following sonication (30 W, 10 sec; Fisher Scientific, USA). Growth rate was calculated as $\mu = (\ln N_2 - \ln N_1) / (t_2 - t_1)$, where N₂ and N₁ are the number of cells during the period of exponential growth at time t₂ and t₁.

Analysis of microcystins : High-pressure liquid chromatography (HPLC) was used to identify and quantify the microcystins. For this analysis, 100 ml of culture was filtered through a 0.45- μm GF/C filter (Whatman, USA) and then lyophilized for 24 hrs in a freeze-drier (Labconco, USA). The lyophilized samples were extracted three times with 50 ml of 5% (v/v) acetic acid for 30 min, while being homogenized with an ultrasonicator (Fisher Scientific, USA). The extract was then centrifuged at 4000 × g for 15 min, and the supernatant was applied to a Sep-Pak C18 cartridge (Waters, USA) that was preactivated by washing with 10 ml of 100% methanol and 10 ml of HPLC-grade distilled water. The cartridge column was washed with 20% methanol and bound microcystins were eluted with 10 ml of 0.1% trifluoroacetic acid in methanol. The eluate was evaporated in a freeze drier (Labconco, USA), and the residue was dissolved in 100% methanol. The solution was then separated by HPLC using an Xterra-C₁₈ column (5 μm ; 4.6 × 15 cm; Waters) and a mobile phase of 52% methanol and 48% 0.05 M phosphate buffer (pH 3.0) (v/v) at a flow rate of 1 ml min⁻¹. Microcystins were detected by measuring the absorbance at 210–420 nm using a photodiode array (Waters, USA).

Results and Discussion

Fig. 1 shows the effects of nitrogen concentration on the growth rate and microcystin production (mg g⁻¹ d.wt.) in *M. aeruginosa* YC, *M. ichthyoblabe* AK and *M. viridis* AK. All three species had maximal growth rates and maximal microcystin production at nitrogen concentrations of 10 to 20 mg l⁻¹.

The specific types of microcystin produced varied according to species and nitrogen concentration. Thus, *M. aeruginosa* YC produced microcystin-YR (MC-YR) when grown in 1-20 mg l⁻¹ nitrogen, but also produced microcystin-RR (MC-RR) and microcystin-LR (MC-LR) when grown in 10-20 mg l⁻¹ N. *M. ichthyoblabe* AK produced MC-YR when grown in 0-10 mg l⁻¹ nitrogen, but produced MC-RR and MC-LR when grown in 20 mg l⁻¹ nitrogen. *M. viridis* AK produced mostly MC-RR and MC-LR when grown in 3-20 mg l⁻¹ nitrogen but produced very little MC-YR at these nitrogen concentrations.

Fig. 2 shows the effects of P concentration on the growth rate and microcystin production of these three *Microcystis*




Fig. 1 : Effect of nitrogen on growth rate (top) and microcystin production (bottom) in three *Microcystis* morphospecies

species. *M. aeruginosa* YC had maximal growth rate and microcystin production at 0.5 mg l⁻¹. *M. ichthyoblabe* AK had maximal growth rate at 0.1 mg l⁻¹ phosphorus and growth rate decreased gradually as phosphorus concentration increased to 5 mg l⁻¹. In contrast, *M. ichthyoblabe* AK produced maximal microcystin at 0 mg l⁻¹ and 5 mg l⁻¹ phosphorus, but minimal microcystin at 0.1 mg l⁻¹ phosphorus. Thus, *M. ichthyoblabe* AK produced more microcystin at phosphorus concentrations that led to lower growth rate.

The growth rate of *M. viridis* AK increased with increasing phosphorus concentration up to 3 mg l⁻¹ and then decreased slightly; its maximum microcystin production was at 0 mg l⁻¹ and 5

mg l⁻¹ phosphorus (as with *M. ichthyoblabe* AK). Interestingly, although this species did not grow at 0 mg l⁻¹ phosphorus, it produced abundant microcystin under this condition. *M. aeruginosa* YC produced all three types of microcystin (MC-RR, MC-YR and MC-LR), but *M. ichthyoblabe* and *M. viridis* only produced two detectable microcystins (MC-RR and MC-LR). *M. aeruginosa* YC produced MC-RR at all tested phosphorus concentrations, and produced the most MC-RR when grown in 0.5 to 5 mg l⁻¹ P. *M. ichthyoblabe* AK also produced MC-RR at all tested phosphorus concentrations, but the concentration of MC-LR was only significant at 5 mg l⁻¹ P. *M. viridis* AK produced MC-RR at all tested concentrations, but produced nearly similar amount of MC-LR at all tested concentrations.

